

A COMPREHENSIVE REVIEW ON MIGRAINE DRUG THERAPY

1. Debjit bhowmik*, Chiranjib, 2. Margret chandira, B. Jayakar

Rajiv Gandhi college of Pharmacy
Nautanwa, Maharajganj, Uttar Pradesh
Vinayaka missions University, Salem, Tamilnadu
Email-debjit_cr@yahoo.com

ABSTRACT

Migraine headaches result from a combination of blood vessel enlargement and the release of chemicals from nerve fibers that coil around these blood vessels. During the headache, an artery enlarges that is located on the outside of the skull just under the skin of the temple (temporal artery). This causes a release of chemicals that cause inflammation, pain, and further enlargement of the artery. A migraine headache causes the sympathetic nervous system to respond with feelings of nausea, diarrhea, and vomiting. This response also delays the emptying of the stomach into the small intestine (affecting food absorption), decreases blood circulation (leading to cold hands and feet), and increases sensitivity to light and sound. Migraine treatment (abortive therapies) and prevention (prophylactic therapies) focus on avoiding triggers, controlling symptoms, and taking medicines. Over-the-counter medications such as naproxen, ibuprofen, acetaminophen (paracetamol), and other analgesics like Excedrin (aspirin with caffeine) are often the first abortive therapies to eliminate the headache or substantially reduce pain. Anti-emetics may also be employed to control symptoms such as nausea and vomiting. Serotonin agonists such as sumatriptan may also be prescribed for severe migraines or for migraines that are not responding to the over-the-counter medications. Similarly, some selective serotonin reuptake inhibitors (SSRIs) - antidepressants such as tricyclics - are prescribed to reduce migraine symptoms although they are not approved in all countries for this purpose. Another class of abortive treatments are called ergots, which are usually effective if administered at the first sign of migraine. Other drugs have also been used to treat migraine such as combinations of barbituates, paracetamol or aspirin, and caffeine (Fioricet or Fiorinal) and combinations of acetaminophen, dichloralphenazone, and isometheptene (Amidrine, Duadrin, and Midrin). If vomiting makes drugs difficult to ingest, anti-emetics will be prescribed. Migraine-specific abortive medications usually are necessary for moderate to severe migraine headaches. The abortive medications for moderate or severe migraine headaches are different than OTC analgesics. Instead of relieving pain, they abort headaches by counteracting the cause of the headache, dilation of the temporal arteries. In fact, they cause narrowing of the arteries. Examples of migraine-specific abortive medications are the triptans and ergot preparations.

INTRODUCTION

A migraine headache is a form of vascular headache. Migraine headache is caused by a combination of vasodilatation (enlargement of blood vessels) and the release of chemicals from nerve fibers that coil around the blood vessels. During a migraine attack, the temporal artery enlarges. (The temporal artery is an artery that lies on the outside of the skull just under the skin of the temple.) Enlargement of the temporal artery stretches the nerves that coil around the artery and causes the nerves to release chemicals. The chemicals cause inflammation, pain, and further enlargement of the artery. The increasing enlargement of the artery magnifies the pain. Migraine attacks commonly activate the sympathetic nervous system in the body. The sympathetic nervous system is often thought of as the part of the nervous system that controls primitive responses to stress and pain, the so-called "fight or flight" response. The increased sympathetic nervous activity in the intestine causes nausea, vomiting, and diarrhea. Sympathetic activity also delays emptying of the stomach into the small intestine and thereby prevents oral medications from entering the intestine and being absorbed. The impaired absorption of oral medications is a common reason for the ineffectiveness of medications taken to treat migraine headaches. The increased sympathetic activity also decreases the circulation of blood, and this leads to pallor of the skin as well as cold hands and feet. The increased sympathetic activity also contributes to the sensitivity to light and sound sensitivity as well as blurred vision. A migraine is a very painful type of headache. People who get migraines often describe the pain as pulsing or throbbing in one area of the head. During migraines, people are very sensitive to light and sound. They may also become nauseated and vomit. Migraine is three times more common in women than in men. Some people can tell when they are about to have a migraine because they see flashing lights or zigzag lines or they temporarily lose their vision. Migraine headaches are usually diagnosed when the symptoms described above are present. Migraine generally begins in childhood to early adulthood. While migraines can first occur in an individual beyond the age of fifty, advancing age makes other types of headaches more likely. A family history is usually present, suggesting a genetic predisposition in migraine sufferers. In addition to diagnosing migraine from the clinical presentation there is usually an accompanying normal examination. Patients with the first headache ever, worst headache ever, or where there is a significant change in headache or the presence of nervous system symptoms, like visual or hearing or sensory loss, may require additional tests. The tests may include blood testing, brain scanning (either CT or MRI), and a spinal tap. Individuals with occasional mild migraine headaches that do not interfere with daily activities usually medicate themselves with over-the-counter (OTC, non-prescription) pain relievers (analgesics). Many OTC analgesics are available. OTC analgesics have been shown to be safe and effective for short-term relief of headache (as well as muscle aches, pains, menstrual cramps, and fever) when used according to the instructions on their labels. There are two major classes of OTC analgesics: acetaminophen (Tylenol) and non-steroidal anti-inflammatory drugs (NSAIDs). The two types of NSAIDs are aspirin and non-aspirin. Examples of non-aspirin NSAIDs are ibuprofen (Advil, Nuprin, Motrin IB, and Medipren) and naproxen (Aleve). Some NSAIDs are available by prescription only. Prescription NSAIDs are usually prescribed to treat arthritis and other inflammatory conditions such as bursitis, tendonitis, etc. The difference between OTC and

prescription NSAIDs may only be the amount of the active ingredient contained in each pill. For example, OTC naproxen (Aleve) contains 220 mg of naproxen per pill, whereas prescription naproxen (Naprosyn) contains 375 or 500 mg of naproxen per pill.

SYMPTOMS OF MIGRAINE

Following are the major **migraine symptoms**:

1. Photophobia
2. Throbbing pain in half of head
3. Eyes become red and patient perceives of burning eyes
4. Nausea sensation and vomiting
5. Loss of appetite
6. Patient wishes to stay all alone during and feels comfortable in silent and dark room
7. Depression and irritability
8. Numbness or weakness in an arm or leg

CAUSES OF MIGRAINE

Following are the major **migraine causes**:

1. Stress and over burdening of mind
2. Tensions
3. Acidity, Indigestion or Constipation
4. Excessive smoking and taking of alcohol
5. Low blood sugar, low blood pressure
6. General body weakness
7. Menstruation in women
8. Nutritional deficiency
9. Consistent overwork
10. Improper sleep and rest.

MIGRAINE HEADACHE DIAGNOSED

Migraine headaches are usually diagnosed when the symptoms described above are present. Migraine generally begins in childhood to early adulthood. While migraines can first occur in an individual beyond the age of fifty, advancing age makes other types of headaches more likely. A family history is usually present, suggesting a genetic predisposition in migraine sufferers. In addition to diagnosing migraine from the clinical presentation there is usually an accompanying normal examination. Patients with the first headache ever, worst headache ever, or where there is a significant change in headache or the presence of nervous system symptoms, like visual or hearing or sensory loss, may require additional tests. The tests may include blood testing, brain scanning (either CT or MRI), and a spinal tap.

APPROACHES OF MIGRAINE HEADACHES PREVENTION

Prevention of migraine

There are two ways to prevent migraine headaches: 1) by avoiding factors ("triggers") that cause the headaches, and 2) by preventing headaches with medications (prophylactic medications). Neither of these preventive strategies is 100% effective. The best one can hope for is to reduce the frequency of headaches.

Migraine triggers

A migraine trigger is any factor that causes a headache in individuals who are prone to develop headaches. Only a small proportion of migraine sufferers, however, clearly can identify triggers. Examples of triggers include stress, sleep disturbances, fasting, hormones, bright or flickering lights, odors, cigarette smoke, alcohol, aged cheeses, chocolate, monosodium glutamate, nitrites, aspartame, and caffeine. For some women, the decline in the blood level of estrogen during the onset of menstruation is a trigger for migraine headaches. The interval between exposure to a trigger and the onset of headache varies from hours to two days. Exposure to a trigger does not always lead to a headache. Conversely, avoidance of triggers cannot completely prevent headaches. Different migraine sufferers respond to different triggers, and any one trigger will not induce a headache in every person who has migraine headaches.

Sleep and migraine

Disturbances such as sleep deprivation, too much sleep, poor quality of sleep, and frequent awakening at night are associated with both migraine and tension headaches, whereas improved sleep habits have been shown to reduce the frequency of migraine headaches. Sleep also has been reported to shorten the duration of migraine headaches.

Fasting and migraine

Fasting possibly may precipitate migraine headaches by causing the release of stress-related hormones and lowering blood sugar. Therefore, migraine sufferers should avoid prolonged fasting.

Bright lights and migraine

Bright lights and other high intensity visual stimuli can cause headaches in healthy subjects as well as patients with migraine headaches, but migraine patients seem to have a lower than normal threshold for light-induced pain. Sunlight, television, and flashing lights all have been reported to precipitate migraine headaches.

Caffeine and migraine

Caffeine is contained in many food products (cola, tea, chocolates, coffee) and OTC analgesics. Caffeine in low doses can increase alertness and energy, but caffeine in high doses can cause insomnia, irritability, anxiety, and headaches. The over-use of caffeine-containing analgesics causes rebound headaches. Furthermore, individuals who consume high levels of caffeine regularly are more prone to develop withdrawal headaches when caffeine is stopped abruptly.

Chocolate, wine, tyramine, MSG, nitrites, aspartame and migraine

Chocolate has been reported to cause migraine headaches, but scientific studies have not consistently demonstrated an association between chocolate consumption and headaches. Red wine has been shown to cause migraine headaches in some migraine sufferers, but it is not clear whether white wine also will cause migraine headaches. Tyramine (a chemical found in cheese, wine, beer, dry sausage, and sauerkraut) can precipitate migraine headaches, but there is no evidence that consuming a low-tyramine diet can reduce migraine frequency. Monosodium glutamate (MSG) has been reported to cause headaches, facial flushing, sweating, and palpitations when consumed in high doses on an empty stomach. This phenomenon has been called Chinese restaurant syndrome. Nitrates and nitrites (chemicals found in hotdogs, ham, frankfurters, bacon and sausages) have been reported to cause migraine headaches. Aspartame, a sugar-substitute sweetener found in diet drinks and snacks, has been reported to trigger headaches when used in high doses for prolonged periods.

Female hormones and migraine

Some women who suffer from migraine headaches experience more headaches around the time of their menstrual periods. Other women experience migraine headaches only during the menstrual period. The term "menstrual migraine" is used mainly to describe migraines that occur in women who have almost all of their headaches from two days before to one day after their menstrual periods. Declining levels of estrogen at the onset of menses is likely to be the cause of menstrual migraines. Decreasing levels of estrogen also may be the cause of migraine headaches that develop among users of birth control pills during the week that estrogens are not taken.

MEDICATION OF MIGRAINE SUFFERERS

Individuals with mild and infrequent migraine headaches that do not cause disability may require only OTC analgesics. Individuals who experience several moderate or severe migraine headaches per month or whose headaches do not respond readily to medications should avoid triggers and consider modifications of their life-style. Life-style modifications for migraine sufferers include:

Beta blockers

Beta-blockers are a class of drugs that block the effects of beta-adrenergic substances such as adrenaline (epinephrine). By blocking the effects of adrenaline, beta-blockers relieve stress on the heart by slowing the rate at which the heart beats. Beta-blockers have been used to treat high blood pressure, angina, certain types of tremors, stage fright, and abnormally fast heart beats (palpitations). They also have become important drugs for improving survival after heart attacks. Beta-blockers have been used for many years to prevent migraine headaches. It is not known how beta-blockers prevent migraine headaches. It may be by decreasing prostaglandin production, though it also may be through their effect on serotonin or a direct effect on arteries. The beta-blockers used in preventing migraine headaches include propranolol (Inderal), atenolol (Tenormin), metoprolol (Lopressor, Lopressor LA, Toprol XL), nadolol (Corgard), and timolol (Blocadren). Beta-blockers generally are well-tolerated. They can aggravate breathing difficulties in patients with asthma, chronic bronchitis, or emphysema. In patients who already have slow heart rates (bradycardias) and heart block (defects in electrical conduction within the heart), beta-blockers can cause dangerously slow heartbeats. Beta-blockers can aggravate symptoms of heart failure. Other side effects include drowsiness, diarrhea, constipation, fatigue, decrease in endurance, insomnia, nausea, depression, dreaming, memory loss, impotence.

Tricyclic antidepressants

Tricyclic antidepressants (TCAs) prevent migraine headaches by altering the neurotransmitters, norepinephrine and serotonin, that the nerves of the brain use to communicate with one another. The tricyclic antidepressants that have been used in preventing migraine headaches include amitriptyline (Elavil), nortriptyline (Pamelor, Aventyl), doxepin (Sinequan), imipramine (Tofranil), and protriptyline. The most commonly encountered side effects associated with TCAs are fast heart rate, blurred vision, difficulty urinating, dry mouth, constipation, weight gain or loss, and low blood pressure when standing. TCAs should not be used with drugs that inhibit monoamine oxidase such as isocarboxazid (Marplan), phenelzine (Nardil), tranylcypromine (Parnate), and procarbazine (Matulane), since high fever, convulsions and even death may occur. TCAs are used with caution in patients with seizures, since they can increase the risk of seizures. TCAs also are used with caution in patients with enlargement of the prostate because they can make urination difficult. TCAs can cause elevated pressure in the eyes of some patients with glaucoma. TCAs can cause excessive sedation when used with other medications that slow the brain's processes, such as alcohol, barbiturates, narcotics, and benzodiazepines, e.g. lorazepam (Ativan), diazepam (Valium), temazepam (Restoril), oxazepam (Serax), clonazepam (Klonopin), zolpidem (Ambien). Epinephrine should not be used with amitriptyline, since the combination can cause severe high blood pressure.

Antiserotonin medications

Methysergide (Sansert) prevents migraine headaches by constricting blood vessels and reducing inflammation of the blood vessels. Methylergonovine is related chemically to

methysergide and has a similar mechanism of action. They are not widely used because of their side effects. The most serious side effect of methysergide is retroperitoneal fibrosis (scarring of tissue around the ureters that carry urine from the kidneys to the bladder). Retroperitoneal fibrosis, though rare, can block the ureters and cause backup of urine into the kidneys. Backup of urine into the kidneys can cause back and flank (the side of the body between the ribs and hips) pain and ultimately can lead to kidney failure. Methysergide also has been reported to cause scarring around the lungs that can lead to chest pain, and shortness of breath.

Calcium channel blockers

Calcium channel blockers (CCBs) are a class of drugs that block the entry of calcium into the muscle cells of the heart and the arteries. By blocking the entry of calcium, CCBs reduce contraction of the heart muscle, decrease heart rate, and lower blood pressure. CCBs are used for treating high blood pressure, angina, and abnormal heart rhythms (e.g., atrial fibrillation). CCBs also appear to block a chemical within nerves, called serotonin, and have been used occasionally to prevent migraine headaches. The CCBs used in preventing migraine headaches are diltiazem (Cardizem, Dilacor, Tiazac), verapamil (Calan, Verelan, Isoptin), and nimodipine.

The most common side effects of CCBs are constipation, nausea, headache, rash, edema (swelling of the legs with fluid), low blood pressure, drowsiness, and dizziness. When diltiazem or verapamil are given to individuals with heart failure, symptoms of heart failure may worsen because these drugs reduce the ability of the heart to pump blood. Verapamil and diltiazem may reduce the elimination and increase the blood levels of carbamazepine (Tegretol), simvastatin (Zocor), atorvastatin (Lipitor), and lovastatin (Mevacor). This can lead to toxicity from these drugs.

Anticonvulsants

Anticonvulsants (antiseizure medications) also have been used to prevent migraine headaches. Examples of anticonvulsants that have been used are valproic acid, phenobarbital, gabapentin, and topiramate. It is not known how anticonvulsants work to prevent migraine headaches.

Prophylactic medications to prevent migraine headaches

Not all migraine sufferers need prophylactic medications; individuals with mild or infrequent headaches that respond readily to abortive medications do not need prophylactic medications. Individuals who should consider prophylactic medications are those who:

- ❖ Require abortive medications for migraine headaches more frequently than twice weekly.
- ❖ Have two or more migraine headaches a month that do not respond readily to abortive medications.

- ❖ Have migraine headaches that are interfering substantially with their quality of life and work.
- ❖ Cannot take abortive medications because of heart disease, stroke, or pregnancy, or cannot tolerate abortive medications because of side effects.

Effective for prophylactic medications

Prophylactic medications can reduce the frequency and duration of migraine headaches but cannot be expected to eliminate migraine headaches completely. The success rate of most prophylactic medications is approximately 50%. Success in preventing migraine headaches is defined as more than a 50% reduction in the frequency of headaches. Prophylactic medications usually are begun at a low dose that is increased slowly in order to minimize side effects. Individuals may not notice a reduction in the frequency, severity, or duration of their headaches for 2-3 months after starting treatment.

Proper way to use preventive medications

Doctors familiar with the treatment of migraine headaches should prescribe preventive medications. Decisions about which preventive medication to use are based on the side effects of the medication and the medical conditions that the patient may have. Propranolol (Inderal) often is used first, provided that the patient does not have asthma, COPD or heart disease. Amitriptyline (Elavil) also is used commonly. Preventive medications are begun at low doses and gradually increased to higher doses if needed. This minimizes side effects from the medications. Preventive medications are to be taken daily for months to years. When they are stopped, the dose needs to be gradually reduced rather than abruptly stopped. Abruptly stopping preventive medications can lead to headaches. In some instances, more than one drug may be needed. Non-medication and behavioral therapies also may be needed.

TREATMENT OF MIGRAINE

- ❖ Avoid factors that cause a migraine attack (for example, lack of sleep, fatigue, stress, certain foods, vasodilators).
- ❖ Treat accompanying conditions (for example, anxiety, depression).
- ❖ Oral birth control agents (contraceptives) may increase the frequency of headaches in females. Women may be advised to discontinue oral contraceptives (or to use a different form) for a trial period to see if they are a factor.

Migraine Headaches, Abortive treatment

Abortive treatments stop migraines quickly. Many drugs are now available for immediate treatment of migraine attacks. The goal is rapid and effective relief of headache. The most effective drugs for stopping a migraine are the triptans, which specifically target serotonin receptors. They are all very similar in chemical structure and action. The following is a list of triptans:

1. Sumatriptan (Imitrex, Imigran)
2. Zolmitriptan (Zomig, Zomig-ZMT)
3. Naratriptan (Amerge, Naramig)
4. Rizatriptan (Maxalt, Maxalt-MLT)
5. Almotriptan (Axert)
6. Frovatriptan (Frova)
7. Eletriptan (Relpax)

The following nontriptans also act on the serotonin receptors. They also act on some other receptors, most likely on those for dopamine and noradrenalin. Sometimes, they are effective when the triptans fail.

1. Ergotamine tartrate (Cafergot)
2. Dihydroergotamine (D.H.E. 45 Injection, Migranal Nasal Spray)
3. Acetaminophen-isometheptene-dichloralphenazone (Midrin)

The following are primarily used when nausea is a complicating factor in migraine headache. In some cases, they also help relieve the headache.

1. Prochlorperazine (Compazine)
2. Promethazine (Phenergan)

Combination drugs like butalbital-acetaminophen-caffeine (Fioricet), butalbital-aspirin-caffeine (Fiorinal), or acetaminophen with codeine (Tylenol With Codeine) are general painkillers in the narcotic class. They can help relieve any kind of pain to some degree, whereas the triptans, ergotamines, and Midrin are used specifically for headaches and do not help relieve arthritis, back pain, or menstrual cramps.

Treatment strategies are more successful if they are tailored to the individual patient and are initiated early in the headache. Patients with severe nausea and vomiting at the onset of an attack may at first respond best to intravenous prochlorperazine. These patients may be

dehydrated; adequate fluid intake is necessary. Vasoconstrictors (agents that narrow the blood vessels), such as ergotamines or triptans, should not be given to patients with known complicated migraine without the advice of a headache specialist. Instead, acute attacks should be treated with one of the other available agents, such as nonsteroidal anti-inflammatory drugs (NSAIDs) or prochlorperazine. Mild and infrequent attacks may not always require the use of ergotamines or triptans and may be adequately treated with acetaminophen (Tylenol), NSAIDs, propoxyphene (Darvon, Darvocet), or a combination of these. About 40% of all attacks do not respond to triptans or any other substance. If all else fails, migraineurs with an attack lasting more than 72 hours (status migrainous) can be treated with intravenous medications. Brief hospitalization may be needed.

MIGRAINE HEADACHES, PREVENTIVE TREATMENT

Patients who have frequent acute migraine attacks and report that the attacks affect their quality of life should consider preventive therapy as a supplement to the specific headache-stopping drugs (abortive treatments) they use. The goals of preventive therapy include decreasing the frequency and severity of acute attacks and improving quality of life. Patients with complicated migraine headaches who have a history of neurological symptoms associated with their attacks are definite candidates for preventive therapy. For these patients, even a single previous complicated migraine episode qualifies them for long-term preventive therapy. The choice of preventive medication should be tailored to the individual's profile, taking into account comorbidities (concurrent medical conditions) such as depression, weight gain issues, exercise tolerance, asthma, and pregnancy plans. All medications have side effects; therefore, selection must be individualized. Preventive drugs include beta-blockers, tricyclic antidepressants, some anticonvulsants, calcium channel blockers, cyproheptadine (Periactin), and NSAIDs such as naproxen (Naprosyn). Unlike the specific headache-stopping drugs (abortive drugs), most of these were developed for other conditions and have been coincidentally found to have headache preventive effects. The following drugs also have preventive effects; unfortunately, they also have more side effects:

- ❖ Methysergide maleate (Sansert): This drug has many side effects.
- ❖ Lithium (Eskalith, Lithobid): This drug has many side effects.
- ❖ Indomethacin (Indocin): This drug can cause psychosis in some people with cluster headaches.
- ❖ Steroids: Prednisone (Deltasone, Meticorten) works extremely well for some people and should be tried if other therapies fail.

How long a person should follow a preventive therapy plan is a function of his or her response to the drug being taken. If headaches completely stop, it is reasonable to gradually reduce the dosage so long as headaches do not recur.

Cluster Headaches

Cluster headaches have been called histamine cephalalgia, Horton neuralgia, and erythromelalgia. The causes of cluster headaches are not known with certainty. The

mechanisms by which the body produces cluster headache pain and other symptoms are also not known for sure.

Cluster Headaches, Prevalence

Cluster headaches are rare. (Less than 1% of the population experience them.) People who do have such headaches usually start to have them when aged 20-40 years. Males get them more often than females (by a ratio of 5-8:1). Usually, no family history of cluster headaches is noted.

Cluster Headaches, Clinical features

Typically, cluster headaches come on with no warning. The signs and symptoms may include intense burning or penetrating pain, often described as a stabbing or hot poker sensation, in or around one eye or temple, occasionally spreading to the forehead, nose, cheek, or upper gum and jaw.

Cluster headaches usually occur on one side of the head. Pain is often penetrating and lasts from 15 minutes to 4 hours. Cluster headaches often cause people to awaken in the middle of the night. During a cluster headache, people are restless and may find relief in pacing or crying. Cluster headaches start rapidly over a few minutes. Periodicity (occurring at regular intervals) is characteristic of cluster headaches. Clusters of headaches are experienced, each cluster lasting as long as several months, once or twice a year. Using alcohol, histamines, or nitroglycerine during a cluster headache may worsen the attack.

Certain personality and physical characteristics have been associated with cluster headaches. A leonine (lionlike) appearance is one of them. Strong associations with smoking, alcohol use, and previous head and face trauma have been noted.

Cluster Headaches, Abortive treatment

Most of the headache-stopping drugs (abortive drugs) effective in treating migraine headaches are also effective in stopping cluster headaches, suggesting that the two types are related.

- Oxygen therapy: This is the treatment of choice and is very safe and effective. Early in an attack, oxygen delivered through a face mask has been known to either stop an attack or diminish its intensity. Why this works is unknown.
- Occipital nerve steroid injection (methylprednisolone acetate [Depo-Medrol]): An injection of this drug may stop a cluster headache attack.

Cluster Headaches, Preventive therapy

As with the abortive drugs, most of the preventive drugs effective in treating migraine headaches are also effective in preventing cluster headaches, again suggesting that the two types are related.

Daily Chronic Headache

Daily chronic headache is defined as a headache that is present for more than 15 days a month and for at least 6 months a year. Three main types are noted: chronic tension-type

headache, migraine chronic tension-type headache complex, and rebound (analgesic abuse) headache. How the body produces chronic daily headaches is not well understood. They have been associated with depression, anxiety, bipolar disorders, panic attacks, mouth/jaw problems, stress, and drug overuse.

Chronic tension-type headache

Chronic tension-type headaches are not associated with a history of migraine or cluster headaches. Patients report almost constant daily headaches of mild-to-moderate intensity. The headache is described as a feeling of tightness or pressure that is not worsened, and may actually be improved, by activity. Patients with chronic tension-type headaches can carry on their daily activities. Nausea and photophobia (sensitivity to light) may occur, but vomiting usually does not. A small group of patients may have head and neck tenderness.

Chronic tension-type headache, Treatment

Patients who are less responsive to previous treatment and those with conditions like depression and stress may be good candidates for psychological treatments. Biofeedback has been successful in patients with tension headache. They are taught how to relax their tense muscles. Thermal biofeedback, in which patients are taught to increase their body temperatures to improve their headaches, has also worked. Other less conventional treatments, such as relaxation training and stress-coping training, may be helpful in the long term.

Transformed migraine

Migraine transformation has been a term used by some experts to describe when intermittent migraines become daily migraines. This type of headache is believed to be associated with analgesic or ergotamine overuse. Patients report intermittent typical migraine attacks along with the daily chronic headaches.

Transformed migraine, Treatment

Stopping all analgesics and headache-related medications is best done in an inpatient setting. Doctors may prescribe a clonidine (Catapres) patch to lessen withdrawal symptoms if narcotic analgesics are involved. Preventive: Preventive treatments for transformed migraine headaches are identical to those used for the other types of migraine headache.

Other uncommon chronic headaches

Hemicrania continua and chronic paroxysmal hemicrania are uncommon forms of chronic headache. Chronic paroxysmal hemicrania is a severe chronic headache similar to cluster headache. It has a male predominance. The headaches are paroxysmal (pulsing), with pain in the temple/eye region lasting 20-30 minutes. The paroxysms occur several times a day. This type of headache can last several years. Treatment with indomethacin (Indocin) results in a dramatic response.

Secondary Headaches

Secondary headaches are related to physical problems and include the following:

- ❖ Space-occupying intracranial (inside the head) lesions: The headaches associated with intracranial tumors are initially paroxysmal. Classic headaches of this type wake a person from sleep at night and are associated with projectile vomiting. With time, the headaches may become continuous and intensify with activities that increase intracranial pressure (for example, coughing, sneezing).
- ❖ Meningeal irritation: Meningitis, especially the chronic forms (tuberculous, fungal), can irritate the meninges (membrane covering the brain and spinal cord) and result in chronic headaches. The headaches are often diffuse (spread out).
- ❖ Posttraumatic headache: Headache can be part of a postconcussion syndrome. Patients may report vague headaches, fatigue, memory problems, and irritability for months or years after the traumatic event.
- ❖ Temporal arteritis: This is an inflammation of some of the arteries of the extracranial (outside the skull) arteries. The headache is generally localized to the affected side and may be worsened by chewing.
- ❖ Post-lumbar puncture (spinal tap) headache: Lumbar puncture can cause a headache that is worsened by sitting up from a lying position. It usually goes away by itself after the person drinks fluids and has caffeine in some form.
- ❖ Referred pain: Headache may be a form of referred pain from neighboring structures. Dental disease can cause chronic headaches. Upper neck diseases or arthritis can also cause headaches. People with acute sinus or jaw problems can experience headaches; however, uncomplicated chronic sinusitis does not cause headaches.
- ❖ Idiopathic intracranial hypertension (benign intracranial hypertension, pseudotumor cerebri): This disorder, most common in young women, is due to increased intracranial (within the head) pressure in the absence of any structural central nervous system abnormality or obstruction to the flow of cerebrospinal fluid.

CONCLUSION

Migraine is often under-diagnosed and under-treated. There is no cure for migraine. Nevertheless, there are numerous interventions that may help restore an improved life for migraine sufferers. These measures should consider the various aspects of the particular patient's condition. Triggering factors, nerve inflammation, blood vessel changes and pain are each addressed aggressively. Individualizing treatment is essential for optimal outcome. The most effective treatment is to identify and avoid the trigger-factors. Aspirin and other analgesics can be taken. Soluble aspirin is better as it is absorbed faster than others and hence if taken at the first sign of headache the severity of the attack is reduced considerably. Many newer drugs are available for the treatment of migraine but are costly and should be kept in reserve. These must be on prescription by the doctor. Alternative therapies like acupuncture, improving posture, stress management, massage of the tensed muscles, etc. are also effective in some cases. This consists of the treatment of any identified underlying cause or disease. The application of cold or heat compresses is done to relieve symptoms. Elimination of food or environmental allergens will help to prevent headaches caused by these factors. Counselling and psychological treatment may also be required. Stress management and biofeedback will also be helpful. Migraine headache is

thought to be due to vascular disturbances. Cluster headaches are also vascular in nature. The exact cause of migraine has not been clearly defined. Vascular disturbances can also be caused by exposure to toxic chemicals like alcohol, lead, arsenic, and carbon monoxide, and are also causes of headache. The most common type of headache is tension headache, but their precise cause is not well-defined. However, most are related to muscle tension, minor trauma, increased stress or anxiety, food and environmental allergens, infection or lesions of the oral or nasal cavity, ear infections, or eyestrain.

REFERENCES:

1. Stephen D. Silberstein, MD, FACP. Practice parameter: Evidence-based guidelines for migraine headache (an evidence-based review): Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology*. 2000;55:754-762.
2. Roger Cady, MD, David W. Dodick, MD. Diagnosis and Treatment of Migraine. *Mayo Clin Proc*. 2002;77:255-261.
3. Dowson AJ, Lipscombe S, Sender J, Rees T, Watson D. New Guidelines for the Management of Migraine in Primary Care. *Curr Med Res Opin*. 2002;18(7):414-439.
4. Patwardhan MB, Samsa GP, Lipton RB, Matchar DB. Changing physician knowledge, attitudes, and beliefs about migraine: evaluation of a new educational intervention. *Headache*. 2006 May;46(5):732-41.
5. Holroyd KA, Drew JB. Behavioral approaches to the treatment of migraine. *Semin Neurol*. 2006 Apr;26(2):199-207.
6. Ramadan NM. Migraine headache prophylaxis: current options and advances on the horizon. *Curr Neurol Neurosci Rep*. 2006 Mar;6(2):95-9.
7. National Guideline Clearinghouse. Treatment of primary headache: acute migraine treatment. Standards of care for headache diagnosis and treatment.

From: Landy S, Smith T. Treatment of primary headache: acute migraine treatment. In: Standards of care for headache diagnosis and treatment. Chicago (IL): National Headache Foundation; 2004. p. 27-39. [11 references].
8. Vincenza Snow, MD. Acute Migraine Treatment Guideline. *Annals of Internal Medicine*. 2003 Oct 1; 139(7):603-4.
9. National Guideline Clearinghouse. Pharmacologic management of acute attacks of migraine and prevention of migraine headache. From: Snow V, Weiss K, Wall EM, Mottur-Pilson C. Pharmacologic management of acute attacks of migraine and prevention of migraine headache. *Ann Intern Med* 2002 Nov 19;137(10):840-52. [121 references].

10. Goetz CG, Pappert EJ. Textbook of Clinical Neurology. 2nd ed. Philadelphia, PA: Saunders; 2003.